

DO-TH 2005/04
astro-ph/0503442
March 2005

Alternative Multipole Vectors for the CMB Temperature Fluctuations

M. Glück and C. Pisano

*Universität Dortmund, Institut für Physik,
D-44221 Dortmund, Germany*

Abstract

We introduce new, low- ℓ , multipole vectors for the CMB temperature fluctuations. A very strong alignment of the quadrupole and octopole vectors is observed as well as a remarkably low declination of these vectors with respect to the galactic plane.

The temperature fluctuations, $\Delta T(\Omega)$, of the CMB radiation are traditionally decomposed according to

$$\Delta T(\Omega) = \sum_{\ell m} a_{\ell m} Y_{\ell m}(\Omega) \quad (1)$$

where $a_{\ell-m} = (-1)^m a_{\ell m}^*$ reflects the reality of $\Delta T(\Omega)$. For each ℓ there are thus $2\ell + 1$ independent real parameters which can be utilized in general to construct ℓ uncorrelated unit vectors and one common overall amplitude.

In [1] a method to construct such multipole vectors was introduced and the properties of these vectors were studied in [1, 2]. In particular, it turned out that the $\ell = 2$ and 3 multipole vectors possess quite unusual correlations which are unexpected for a statistically isotropic sky with Gaussian random $a_{\ell m}$, $\langle a_{\ell m} a_{\ell' m'}^* \rangle = C_\ell \delta_{\ell\ell'} \delta_{mm'}$. The multipole vectors presented in [1] were obtained by the construction of symmetric traceless rank- ℓ tensor representation of the spherical harmonics $Y_{\ell m}$ and it remains to be seen to what extent the above mentioned correlations depend on these particularly chosen multipole vectors.

To study this question we introduce an alternative set of multipole vectors whose directions point towards the maxima or minima of $\Delta T_\ell(\Omega) \equiv \sum_m a_{\ell m} Y_{\ell m}(\Omega)$. Utilizing the $a_{\ell m}$ in Table II of [3], obtained from a foreground cleaned [4] CMB map one obtains in galactic coordinates (l, b) the following unit vectors:

$$\begin{aligned} \hat{n}_1 &= (157.03^\circ, 6.29^\circ), \\ \hat{n}_2 &= (63.33^\circ, 30.41^\circ), \end{aligned} \quad (2)$$

in the quadrupole sector and

$$\begin{aligned} \hat{n}_a &= (145.97^\circ, 10.59^\circ), \\ \hat{n}_b &= (14.59^\circ, 29.75^\circ), \\ \hat{n}_c &= (82.09^\circ, 17.93^\circ), \end{aligned} \quad (3)$$

in the octopole sector. For convenience these vectors were chosen to point towards the north galactic hemisphere. We note the rather unusual alignments between some of the $\ell = 2$ and $\ell = 3$ vectors, i.e.:

$$\hat{\mathbf{n}}_1 \cdot \hat{\mathbf{n}}_a = 0.9787, \quad \hat{\mathbf{n}}_2 \cdot \hat{\mathbf{n}}_c = 0.9328, \quad (4)$$

which in a statistically isotropic sky with random $a_{\ell m}$ is expected to happen by chance only about once in $(1 - \hat{\mathbf{n}}_1 \cdot \hat{\mathbf{n}}_a)^{-1}(1 - \hat{\mathbf{n}}_2 \cdot \hat{\mathbf{n}}_c)^{-1} \simeq 700$, (cf. footnote 4 in [3]). Similar alignments also exist for the multipole vectors $\hat{\mathbf{v}}^{(\ell,i)}$ of [2]:

$$\hat{\mathbf{v}}^{(2,1)} \cdot \hat{\mathbf{v}}^{(3,1)} = 0.9730, \quad \hat{\mathbf{v}}^{(2,2)} \cdot \hat{\mathbf{v}}^{(3,2)} = 0.8636, \quad (5)$$

which is expected to happen by chance only about once in 270, quite lower than for the previous alignments. The coplanarity of our unit vectors is characterized by

$$D_{ab} = 0.9715, \quad D_{ac} = 0.9693, \quad D_{bc} = 0.9007, \quad (6)$$

where, following [1, 2], $D_{ab} \equiv |(\hat{\mathbf{n}}_1 \times \hat{\mathbf{n}}_2) \cdot (\hat{\mathbf{n}}_a \times \hat{\mathbf{n}}_b)| / |\hat{\mathbf{n}}_1 \times \hat{\mathbf{n}}_2| |\hat{\mathbf{n}}_a \times \hat{\mathbf{n}}_b|$, with analogous definitions for D_{ac} and D_{bc} . The corresponding products for the vectors $\hat{\mathbf{v}}^{(\ell,i)}$ are [2]:

$$D_1 = 0.9531, \quad D_2 = 0.8719, \quad D_3 = 0.8377, \quad (7)$$

i.e. a somewhat reduced coplanarity between the $\ell = 2$ and $\ell = 3$ vectors as compared to (6). This reduction is directly related to the corresponding reduced alignments in (5) as compared to (4).

Finally we note the lower declinations $b(\hat{\mathbf{n}})$ as compared to $b(\hat{\mathbf{v}})$ implying a closer vicinity of our multipole vectors to the galactic plane. This close vicinity, characterized by $\cos b(\hat{\mathbf{n}}_1) = 0.9940$, and the apparently unrelated but unusual alignments of our $\ell = 2$ and $\ell = 3$ unit vectors, as given in (4), suggest a common origin for these – seemingly unrelated – remarkable correlations.

To summarize, we have introduced a new set of $\ell = 2, 3$ multipole vectors pointing towards the maxima or minima of the corresponding CMB temperature fluctuations.

These vectors are characterized by an unusual high alignment between some quadrupole and octopole unit vectors as well as by a simultaneous very low declination of these vectors with respect to the galactic plane.

This work has been supported in part by the ‘Bundesministerium für Bildung und Forschung’, Berlin/Bonn.

References

- [1] C. J. Copi, D. Huterer and G. D. Starkman, Phys. Rev. D **70**, 043515 (2004).
- [2] D. J. Schwarz, G. D. Starkman, D. Huterer and C. J. Copi, Phys. Rev. Lett. **93**, 221301 (2004).
- [3] A. de Oliveira-Costa, M. Tegmark, M. Zaldarriaga and A. Hamilton, Phys. Rev. D **69**, 063516 (2004).
- [4] M. Tegmark, A. de Oliveira-Costa and A. Hamilton, Phys. Rev. D **68**, 123523 (2003).